

The Method of Partial Fractions

I. Introduction

Another integration technique is called the *method of partial fractions*. This technique is used to find the antiderivatives for a certain class of functions, the rational functions. Recall, a rational function is a function of the form $f(x) = \frac{p(x)}{q(x)}$ where $p(x)$ and $q(x)$ are polynomial functions. For example:

$\int \frac{6x^3 - 3x^2 + 7x - 1}{2x^2 - x - 1} dx$ has a rational function for its integrand. There is no obvious function whose derivative is this rational function. If we can rewrite such a rational function as the sum of simpler rational functions that are easier to integrate, we have a method for evaluating the integral. Simplifying a rational function to the sum of simpler rational functions is called *partial fraction decomposition*. The general steps to evaluating these types of integrals are:

1. Determine if the integrand is a *proper* rational function ($\deg p(x) < \deg q(x)$). If it is not, use long division to rewrite the integrand as $\frac{p(x)}{q(x)} = Q(x) + \frac{r(x)}{q(x)}$ where $Q(x)$ is a quotient and $r(x)$ a remainder.
2. Factor the denominator, $q(x)$, of the proper rational function and determine the proper form to use for the partial fraction decomposition:

(a) If $q(x)$ contains **unique linear factors** use
$$\frac{p(x)}{q(x)} = \frac{A_1}{a_1x + b_1} + \frac{A_2}{a_2x + b_2} + \dots + \frac{A_n}{a_nx + b_n}$$

(b) If $q(x)$ contains **repeated linear factors** use
$$\frac{p(x)}{q(x)} = \frac{A_1}{a_1x + b_1} + \frac{A_2}{(a_1x + b_1)^2} + \dots + \frac{A_n}{(a_1x + b_1)^n}$$

- (c) If $q(x)$ contains **irreducible quadratic factors** use

$$\frac{p(x)}{q(x)} = \frac{A_1x + B_1}{a_1x^2 + b_1x + c_1} + \frac{A_2x + B_2}{a_2x^2 + b_2x + c_2} + \dots + \frac{A_nx + B_n}{a_nx^2 + b_nx + c_n}$$

- (d) If $q(x)$ contains **repeated quadratic factors** use

$$\frac{p(x)}{q(x)} = \frac{A_1x + B_1}{a_1x^2 + b_1x + c_1} + \frac{A_2x + B_2}{(a_1x^2 + b_1x + c_1)^2} + \dots + \frac{A_nx + B_n}{(a_1x^2 + b_1x + c_1)^n}$$

Note: $q(x)$ is likely to contain mixed factors that will require a combination of these forms.

3. Equate the proper rational function $\frac{p(x)}{q(x)}$ with the form obtained above. Solve for the constants.

4. Rewrite the integrand, $\frac{p(x)}{q(x)}$, in the decomposed form obtained and solve the integral.

Example 1:

Evaluate $\int \frac{10x-1}{2x^2-x-1} dx$

1. Is degree $p(x) <$ degree $q(x)$? (If not, use long division first.)

2. Partial Fraction Decomposition:
 - (a) Factor $q(x)$. Based on the factors obtained, set up the needed form(s).
 - (b) Multiply both sides by the least common denominator.
 - (c) Simplify right side and collect coefficients of like terms.
 - (d) Equate coefficients on left and right sides to form a system.
 - (e) Solve the system for A, B, etc.

3. Integrate the decomposed form of $\frac{p(x)}{q(x)}$.

Example 2: For each rational expression below, set up the proper form for partial fraction decomposition. DO NOT solve for the unknown constants.

(a) $\frac{x}{(x+3)^2}$

(b) $\frac{x-5}{x^3+x^2}$

(c) $\frac{2}{x^3+5x^2+8x}$

(d) $\frac{x^3}{x^2+1}$

Example 3: Solve $\int \frac{10}{(x-1)^2(x^2+9)} dx$

You may use your textbook, lab and notes. Students may work cooperatively but must submit their own set of Lab Exercises. **No calculators.**

A. For each rational function, perform steps 1, 2, and 3 to set up the partial fraction decomposition of the integrand. DO NOT solve for the unknown constants.

1. $\frac{5x^2 - 3x + 2}{x^3 - 4x}$

2. $\frac{1}{x^4 - 10x^2 + 9}$

3. $\frac{x^2}{(x-1)(x^2+1)^2}$

B. For each integral, perform steps 1-4 to evaluate. Show all your work, identify substitutions.

1. $\int \frac{12x-1}{x^2-5x+6} dx$

2. $\int \frac{2x^2 + 7x + 4}{x^2 + 2x} dx$

3. $\int \frac{10}{(x-1)^2(x^2+9)} dx$